

(ofc)	971.801.3394 (cell)	503.339.9568 (fax)	extrationpractice@gmail.com
	www.Ore	egonWisdomTeeth.com	L

PRE-OPERATIVE INSTRUCTIONS FOR DENTAL SURGERY

** VERY IMPORTANT INFORMATION—PLEASE READ CAREFULLY ** ** COMPLETE ATTACHED "MEDICAL HISTORY UPDATE FORM" ** & RETURN IT TO YOUR DENTIST PRIOR TO SURGERY

- 1. If you have any concerns or questions about the surgery, please contact Dr. Brannen at 971.801.3394 or by email at paul@drbrannen.com.
- 2. I will be reviewing your medical history with you immediately prior to the surgery. Please be sure you are familiar with that information especially with the name(s) and dosage(s) of any medications you are taking. If you feel your history is relatively complicated, we will need to decide if a consultation with your physician is necessary before the procedure is performed.
- 3. Patients who are minors (under 18 years of age) must have a legal guardian present to both fill out the "Medical History Update Form" and to sign the "Disclosure and Consent Form."
- 4. It is important to avoid smoking for at least one week before the surgery and one week following the surgery.
- 5. Keep in mind that it is best to allow for some flexibility around your appointment time on the day of your surgery. It is best not to "squeeze in" an appointment for surgery on an already busy day.

If you are having IV (Intravenous) conscious sedation:

- 1. To reduce the chances of nausea, do not eat or drink anything (including water) for <u>at least six hours</u> prior to your appointment.
 - If your surgery is in the morning, do not eat or drink anything between bedtime and your scheduled appointment.
 - If your surgery is in the afternoon, a light breakfast before 7:00 a.m. is encouraged.
 - Unless specified by your dentist, all medicines taken on a routine basis should be continued without interruption. Please swallow with a minimal amount of water.
- 2. A responsible adult, over 18 years of age, should accompany you to the office and should <u>remain</u> <u>in the office during the entire procedure</u>. Following the sedation, this responsible adult should be physically capable of assisting and accompanying you home and should remain with you for the next 24 hours.
- 3. If receiving intravenous sedation, you should wear clothing, which is not restricting to the neck or arms. You should wear loose-fitting tops on which the sleeves can be rolled up to the shoulder. Also, please be sure to wear shoes that are securely fastened; no flip-flops or loose-fitting sandals, please.
- 4. Following the sedation, you should refrain from driving an automobile or engaging in any activity that requires alertness for the next 24 hours.
- 5. There are important differences between general anesthesia (being completely asleep) and IV conscious sedation. If you have any questions about the IV conscious sedation process, please feel free to contact Dr. Brannen at 971.801.3394 prior to the procedure.

NOTE: Additional pre-operative information can be found at www.drbrannen.com. I recommend you preview the "Disclosure and Consent Form" on the website, or you can request a copy from your dentist.



(ofc) 971.801.3394 (cell) 503.339.9568 (fax) extrationpractice@gmail.com www.OregonWisdomTeeth.com

MEDICAL HISTORY UPDATE FORM

						Date		
Jame_						Dentist's Name:		
	Last	First		Middle	_			
ocial S	Security #	Ht		Wt		Date of Birth		
f you a	are completing this fo	rm for another person, wi	hat is y	our relation	onship to	that person?		
	nfidential. Please not		isit, you	ı will be a	sked some	e for our records only and wil e questions about your respon cerning your health.		
1. 2. 3. 4.	Has there been any che health within the past My last physical exam Are you now under the physician?	year?	No	i. j. k. l. m. n. o.	AIDS or I Thyroid p Respirato Stomach Kidney tr High or lo Sexually Epilepsy/	jaundice, or liver disease HIV infection	Yes Yes Yes Yes Yes Yes Yes	No
6. 7.	hospitalized in the pas Are you taking any monon-prescription media	ous illness, operation, or be t 5 years?	No No	r. 10. Ha Or 11. Do as 12. Ha	Sleep apn ave you had r required a to you have anemia? ave you be	d abnormal bleeding?	Yes Yes Yes Yes	No No No No
8.9.	Do you have or have y diseases or problems? a. Damaged or artific murmur, or rheuma b. Cardiovascular diseattack, heart trouble	Boniva?		a. b. c. d. e. f. g. h.	Local and Penicillin Sulfa drug Barbitura Aspirin Iodine Codeine of Other	rgic or have you had a reaction esthetics	Yes Yes Yes Yes Yes Yes	No No No No No No
have errors would	d. Cancer requiring IV e. Asthma or hay feve f. Fainting spells or s g. Diabetes ify that I have read and been answered to my s s or omissions that I ma d like to provide us wit	atisfaction. I will not hold y have made in the comple h additional information, it	knowled I my der tion of t	15. Do 16. An 17. An ge that my ntist, or an his form.	re you pregot you have re you nurs re you taking questions the regular mounts of the regular mounts regular mounts regular mounts regular regu	any menstrual problems? any menstrual problems? ng birth control pills? s, if any, about the inquiries set ember of his/her staff, responsedical history is complex or if you would use the back of this for	Yes Yes Yes forth ible for	No above or any el you
	chronological narrative	of your medical history.		Sign	ature of Pa	tient (or Patient's Guardian)		





PAUL A. BRANNEN, DMD, LLC EXTRACTION RECORD

Procedure to be completed in the office of	Patient Name	Age	DOB	//_	Date//
Tech to be Extracted: #\$ Fee \$ AF\$ \$F\$ CC " "Pt Phone"	Procedure to be completed in the office of	Pt Er			
Post-Op Ride	Teeth to be Extracted: #'s	Fee \$		_A/F \$	S/F \$
Post-Op Ride	CC"	"Pt Phone # (_)		
Findings/Diagnosis:	Post-Op Ride Po	st-Op Ride's # ()		
Findings/Diagnosis:	Radiograph(s): Pano PA Other Date / /	Exam/Consult:	Y N Ass	sistants	
Rear Pericononist#					
Rear Pericononist#			☐ RCT decl	ined#	
SymptAsympt#	Recur Pericoronitis# Periodontal Disease #		☐ Pre-prosth	etic#	
ASA: I II III Mallampati: I II III IV BMI Informed Consent/PARO: Obtained both written and verbal, patient and/or legal guardian had opportunity to ask questions Pre-Operative Medication BP	☐ Sympt/Asympt# ☐ Irrever/Necro/Prev RCT #	#	Other#		
Informed Consent/PARO: Obtained both written and verbal, patient and/or legal guardian had opportunity to ask questions Pre-Operative Medication BP			_		
Obtained both written and verbal, patient and/or legal guardian had opportunity to ask questions Pre-Operative Medication See Medical History Form See Sedation Record NKDA Allergies					
Pre-Operative Medication Medical Hx: See Medical History Form See Sedation Record NKDA Allergies Local Anesthesia: 2% Lido w/ 1:100k epi, carts 0.5% Mare w/ 1:200k epi, carts 4.5% Edward Medical History Form See Sedation Record NKDA Allergies Local Anesthesia: 2% Lido w/ 1:100k epi, carts 0.5% Mare w/ 1:200k epi, carts 4.5% Edward Medical Microscope Microscope Medical Microscope Microscope Medical Microscope Microscope Medical Microscope Microscope Microscope Medical Microscope Microsco		opportunity to ask	questions		
Medical Hx: See Medical History Form See Sedation Record NKDA Allergies			-	bpn	n
Local Anesthesia: 2% Lido w/ 1:100k epi,carts 0.5% Marc w/ 1:200k epi,carts 4% Septo w/ 1:100k epi,carts 3% Carbo,carts Other					
4% Septo W 1:100k epi, carts 3% Carbo, carts Other	· · · · · · · · · · · · · · · · · · ·				
Procedure: Non-surgical EXT # BB and throat barrier placed; tissue released; elevator & forcep removal whole; curette granulation tissue prn; profuse irrigation w/ 0.9% NaCl solution; damp gauze placed for hemostasis					
Non-surgical EXT # BB and throat barrier placed; tissue released; elevator & forcep removal whole; curette granulation tissue pm; profuse irrigation w/ 0.9% NaCl solution; damp gauze placed for hemostasis Justification					
BB and throat barrier placed; tissue released; elevator & forcep removal whole; curette granulation tissue prn; profuse irrigation w/ 0.9% NaCl solution; damp gauze placed for hemostasis BB and throat barrier placed; FTMPF; peripheral ostectomy; tooth sectioned w/ HP under copious irrigation, tooth removed w/ elevator and forceps; bone file and/or Rongeur used prn; curette granulation tissue prn; profuse irrigation w/ 0.9% NaCl; damp gauze placed for hemostasis Suture(s) placed; Vicryl; Silk; PTFE Allooss (50/50 cort/cane; cancellous; cortical) mixed w/ 0.9% NaCl and placed in socket/defect, RCM6 collagen membrane trimmed and positioned under tissue flap; Suture(s): Gut; Vicryl; Silk; PTFE Other Tooth removed whole w/ elevator and forcep Tooth removed whole w/ elevator and forcep Tooth removed whole w/ elevator and forcep DB incision w/ M vertical release Tooth removed in pieces w/ elevator and forcep Damp gauze place for hemostasis Damp gauze place Gut; Vicryl; Silk; PTFE Tooth removed in pieces w/ elevator and forcep Tooth removed in pieces w/ elevator and forcep Tooth removed whole w/ elevator and forcep Tooth removed in pieces w/ elevator and forcep Tooth removed whole w/ elevator and forcep DB incision w/ M vertical release DB in					
Surgical EXT # Surg		al whole; curette gr	anulation tis	sue prn; pro	fuse irrigation w/
□ Surgical EXT # Justification BB and throat barrier placed; FTMPF; peripheral ostectomy; tooth sectioned w/ HP under copious irrigation, tooth removed w/ elevator and forceps; bone file and/or Rongeur used prn; curette granulation tissue prn; profuse irrigation w/ 0.9% NaCl; damp gauze placed for hemostasis □ Suture(s) placed; Gut; Vicryl; Silk; PTFE □ Allooss (50/50 cort/cane; cancellous; cortical) □ Foundation trimmed and placed; □ Suture(s): Gut; Vicryl; Silk; PTFE □ Allooss (50/50 cort/cane; cancellous; cortical) mixed w/ 0.9% NaCl and placed in socket/defect, RCM6 collagen membrane trimmed and positioned under tissue flap; Suture(s): Gut; Vicryl; Silk; PTFE □ Other Maxillary 3 rd molars # □ Tooth removed whole w/ elevator and forcep □ BB and throat barrier placed □ Tooth removed whole w/ elevator and forcep □ BB incision w/ M vertical release □ Curette granulation tissue prn □ BB incision intrasulcular to 1 rd molar □ Profuse irrigation w/ 0.9% NaCl □ Peripheral ostectomy to fully uncover tooth □ Damp gauze place for hemostasis □ BB, MB, trough □ Tooth removed whole w/ elevator and forcep □ Mandibular 3 rd molars # □ Tooth removed whole w/ elevator and forcep □ TMPF □ FTMPF □ Follicular tissue removed □ No incision □ Tooth removed whole w/ elevator and forcep □ BB and throat barrier placed □ Tooth removed whole w/ elevator and forcep □ DB incision intrasulcular to 1 rd molar □ Curette granulation tissue prn □ Peripheral ostectomy to fully uncover tooth □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes		, 3		1 /1	Ç
elevator and forceps; bone file and/or Rongeur used prn; curette granulation tissue prn; profuse irrigation w/ 0.9% NaCl; damp gauze placed for hemostasis Suture(s) placed: Gut; Vicryl; Silk; PTFE Allooss (50/50 cort/cane; cancellous; cortical) Profundation trimmed and placed; Suture(s): Gut; Vicryl; Silk; PTFE Allooss (50/50 cort/cane; cancellous; cortical) Rongeur w/ 0.9% NaCl and placed in socket/defect, RCM6 collagen membrane trimmed and positioned under tissue flap; Suture(s): Gut; Vicryl; Silk; PTFE Other Tooth removed whole w/ elevator and forcep Pollicular tissue removed Profuse irrigation w/ 0.9% NaCl Peripheral ostectomy to fully uncover tooth Damp gauze place for hemostasis Tooth removed whole w/ elevator and forcep Tooth removed in pieces w/ elevator and forcep Tooth removed in pieces w/ elevator and forcep Tooth removed whole w/ elevator and forcep Dab incision w/ M vertical release Bone file/Rongeur used to smooth bone Dab Bincision intrasulcular to 1st molar Curette granulation tissue prn Profuse irrigation w/ 0.9% NaCl Damp gauze place for hemostasis Incomplete B/L section w/ HP, completed w/ elevator Damp gauze place for hemostasis Damp gau					
placed for hemostasis	BB and throat barrier placed; FTMPF; peripheral ostectomy; tooth sec	tioned w/ HP under	copious irri	gation, tooth	removed w/
Ridge Pres/Bone Graft #	elevator and forceps; bone file and/or Rongeur used prn; curette granu	lation tissue prn; pr	ofuse irrigati	ion w/ 0.9%	NaCl; damp gauze
□ Foundation trimmed and placed;Suture(s): Gut; Vicryl; Silk; PTFE □ Allooss (50/50 cort/canc; cancellous; cortical) mixed w/ 0.9% NaCl and placed in socket/defect, RCM6 collagen membrane trimmed and positioned under tissue flap; Suture(s): Gut; Vicryl; Silk; PTFE □ Other Maxillary 3 rd molars # □ Tooth removed whole w/ elevator and forcep □ No incision □ Follicular tissue removed □ Follicular tissue removed □ DB incision w/ M vertical release □ Curette granulation tissue prn □ DB incision intrasulcular to 1 st molar □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ DB, B, MB, trough □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ Tooth removed in pieces w/ elevator and forcep □ Tooth removed whole w/ elevator and forcep □ DB incision w/ M vertical release □ Tooth removed whole w/ elevator and forcep □ DB incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ Curette granulation tissue prn □ Peripheral ostectomy to fully uncover tooth □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Damp gauze place	placed for hemostasis \(\bigcup_\) Suture(s) placed: Gut; Vicryl; Silk; P	TFE			
mixed w/ 0.9% NaCl and placed in socket/defect, RCM6 collagen membrane trimmed and positioned under tissue flap; Suture(s): Gut; Vicryl; Silk; PTFE Other DB and throat barrier placed	☐ Ridge Pres/Bone Graft # Justification				
Suture(s): Gut; Vicryl; Silk; PTFE Other Maxillary 3 rd molars # BB and throat barrier placed Tooth removed whole w/ elevator and forcep No incision Follicular tissue removed Bone file/Rongeur used to smooth bone Curette granulation tissue prn DB incision intrasulcular to 1 st molar Profuse irrigation w/ 0.9% NaCl Peripheral ostectomy to fully uncover tooth Damp gauze place for hemostasis DB, B, MB, trough Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # BB and throat barrier placed Tooth removed in pieces w/ elevator and forcep No incision Tooth removed whole w/ elevator and forcep TTMPF Follicular tissue removed DB incision intrasulcular to 1 st molar Follicular tissue removed DB incision intrasulcular to 1 st molar Curette granulation tissue prn Peripheral ostectomy to fully uncover tooth Profuse irrigation w/ 0.9% NaCl DB, B, MB, trough Damp gauze place for hemostasis Incomplete B/L section w/ HP, completed w/ elevator Damp gauze place for hemostasis DB incision intrasulcular to 1 st molar Curette granulation tissue prn Profuse irrigation w/ 0.9% NaCl DB map gauze place for hemostasis DB map gauze pla					
Bal and throat barrier placed Tooth removed whole w/ elevator and forcep Policular tissue removed Bone file/Rongeur used to smooth bone DB incision w/ M vertical release Curette granulation tissue pm DB incision intrasulcular to 1st molar Profuse irrigation w/ 0.9% NaCl Peripheral ostectomy to fully uncover tooth Damp gauze place for hemostasis DB, B, MB, trough Suture(s) placed: Gut; Vicryl; Silk; PTFE Tooth removed in pieces w/ elevator and forcep Tooth removed in pieces w/ elevator and forcep Tooth removed whole w/ elevator and forcep Tooth removed whole w/ elevator and forcep DB incision w/ M vertical release Bone file/Rongeur used to smooth bone DB incision intrasulcular to 1st molar Curette granulation tissue prn Peripheral ostectomy to fully uncover tooth Damp gauze place for hemostasis Incomplete B/L section w/ HP, completed w/ elevator Damp gauze place for hemostasis Incomplete B/L section w/ HP, completed w/ elevator Damp gauze place for hemostasis		nbrane trimmed and	d positioned	under tissue	flap;
□ BB and throat barrier placed □ No incision □ Follicular tissue removed □ FTMPF □ Bone file/Rongeur used to smooth bone □ Curette granulation tissue prm □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # □ BB and throat barrier placed □ No incision □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ BB incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ User Pertinent Notes □ Other Pertinent Notes					
□ No incision □ FTMPF □ Bone file/Rongeur used to smooth bone □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3rd molars # □ BB and throat barrier placed □ No incision □ BB incision w/ M vertical release □ BB incision w/ M vertical release □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Tooth removed in pieces w/ elevator and forcep □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vieryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes					_
□ FTMPF □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3rd molars # □ BB and throat barrier placed □ Tooth removed in pieces w/ elevator and forcep □ No incision □ DB incision w/ M vertical release □ Tooth removed in pieces w/ elevator and forcep □ FTMPF □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □ Profuse irrigation w/ 0.9% NaCl □ Other Pertinent Notes □ Profuse rinse 4oz x 1 □ Norco 5/325mg x 6 □ Other Pertinent Notes □ Curette granulation tissue prn □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6				levator and f	orcep
□ DB incision w/ M vertical release □ Curette granulation tissue prn □ DB incision intrasulcular to 1 st molar □ Profuse irrigation w/ 0.9% NaCl □ Peripheral ostectomy to fully uncover tooth □ Damp gauze place for hemostasis □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ Bone file/Rongeur used to smooth bone □ DB incision intrasulcular to 1 st molar □ Curette granulation tissue prn □ Peripheral ostectomy to fully uncover tooth □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □ Norco 5/325mg x 6					
□ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ Damp gauze place for hemostasis □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3rd molars # □ BB and throat barrier placed □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ BB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Profuse irrigation w/ 0.9% Caut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes					;
□ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # □ BB and throat barrier placed □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ Bi incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes					
□ DB, B, MB, trough □ Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # □ BB and throat barrier placed □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ BB incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes					
□ Tooth removed in pieces w/ elevator and forcep Mandibular 3 rd molars # □ BB and throat barrier placed □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ Bone file/Rongeur used to smooth bone □ DB incision intrasulcular to 1 st molar □ Curette granulation tissue pm □ Peripheral ostectomy to fully uncover tooth □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □					Y'II DEEE
□ BB and throat barrier placed □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ Bone file/Rongeur used to smooth bone □ DB incision intrasulcular to 1st molar □ Curette granulation tissue prn □ Peripheral ostectomy to fully uncover tooth □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes □ Tooth removed in pieces w/ elevator and forcep □ Tooth removed whole w/ elevator and forcep □ Follicular tissue removed □ Bone file/Rongeur used to smooth bone □ Curette granulation tissue prn □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes		Suture(s	s) placed: Gu	it; Vicryl; S	filk; PIFE
□ BB and throat barrier placed □ Tooth removed in pieces w/ elevator and forcep □ No incision □ Tooth removed whole w/ elevator and forcep □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ Bone file/Rongeur used to smooth bone □ DB incision intrasulcular to 1st molar □ Curette granulation tissue prn □ Peripheral ostectomy to fully uncover tooth □ Profuse irrigation w/ 0.9% NaCl □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes □ Tooth removed in pieces w/ elevator and forcep □ Tooth removed whole w/ elevator and forcep □ Follicular tissue removed □ Bone file/Rongeur used to smooth bone □ Curette granulation tissue prn □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes	■ 1 ooth removed in pieces w/ elevator and forcep				
□ No incision □ FTMPF □ Follicular tissue removed □ DB incision w/ M vertical release □ DB incision intrasulcular to 1st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes				/ 1 4	1.0
□ FTMPF □ Bone file/Rongeur used to smooth bone □ DB incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □ Post-Other □ Post-Other Pertinent Notes □ Post-Other Pertinent Notes □ Post-Other □ Post-Other Pertinent Notes □ Post-Other □ Post-Othe					
□ DB incision w/ M vertical release □ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □ Curette granulation tissue prn □ Curette granulation tissue prn □ Curette granulation tissue prn □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ O				ievator and i	orcep
□ DB incision intrasulcular to 1 st molar □ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □ Curette granulation tissue prn □ Curette granulation tissue prn □ Profuse irrigation w/ 0.9% NaCl □ Damp gauze place for hemostasis □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Ry: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □				ama a atla la ama	
□ Peripheral ostectomy to fully uncover tooth □ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other □ Other Pertinent Notes □					;
□ DB, B, MB, trough □ Damp gauze place for hemostasis □ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes					
□ Incomplete B/L section w/ HP, completed w/ elevator □ Suture(s) placed: Gut; Vicryl; Silk; PTFE □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes					
 □ Post-Operative Instructions: Both written and verbal given to patient and/or legal guardian and escort. Extra gauze and Monoject syringe provided. Release signed by escort. Rx: □ Pen VK 500mg x 20 □ Clindamycin 150gm x 20 □ Peridex rinse 4oz x 1 □ Norco 5/325mg x 6 □ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes 					Sile DTEE
Monoject syringe provided. Release signed by escort. Rx: □Pen VK 500mg x 20 □Clindamycin 150gm x 20 □Peridex rinse 4oz x 1 □Norco 5/325mg x 6 □Ibuprofen 600mg x 30 □Other Other Pertinent Notes					
Rx: Pen VK 500mg x 20 Clindamycin 150gm x 20 Peridex rinse 4oz x 1 Norco 5/325mg x 6 Dibuprofen 600mg x 30 Other Other Pertinent Notes		panem and/or legal	guaruian an	u escort. Ext	ia gauze allu
□ Ibuprofen 600mg x 30 □ Other Other Pertinent Notes		nse 4oz v 1 DNon	co 5/325ma	v 6	
Other Pertinent Notes			_	λ 0	
	• • •		_		
Doctor's Signature Date	Other Pertinent Notes				
	Doctor's Signature	ī	Date.		

Patient Name	DOB/(2 of 2)
POST-OPERATIVE RECORD	(= V1 =)
□ Post-Op Call/Text Date:/ □ Pt d	id not answer. Left Message/Unable to leave message
☐ Patient reports they are doing well with swelling and p answered all questions.	ain within normal limits for treatment completed;
☐ Reinforced post-op instructions (i.e., diet, activity, med	dication instructions, contact info, etc.)
☐ Pain Scale (<i>circle</i>) 0 1 2 3 4 5 6 7 8 9 10	
Notes:	
Doctor's Signature	
Additional Notes:	
	☐ Drug Log Recorded// ☐ Posted for Payment / /
	Comment Card Sent/
	☐ 1 Week Post-Op Call/
Doctor's Signature	Date



(ofc)	971.801.3394 (cell)	503.339.9568 (fax)	extrationpractice@gmail.com
	W	ww.OregonWisdom	Teeth.com

MODERATE SEDATION RECORD

Date																n.	e-ap \	/itala		BP			HR				Sp()2			RR	
Name																	200															
																Ag	je	G	ender	٧	Veight		Hei	ght	BI	MI		Malla	mpa	ti	ASA	
M. R I I II - A																1		М	F	:		lbs	ŝ.				_				<u>L</u>	
Medical History																NF	0											NIBP	Ovi	L mete		3
																So	lids		Н	lours	(Clear L	iquid	s		Hours	8	Prec	ordia	l	4	
																	ocedu	re(s)										ECG				
Medications																												EtCO)2			
																Or	perato	r.						Acc	istant	· c•	_		_	—		
																10	oraco							133	is carr							
Drug Allergies																Г		S	tart		Finis	sh		Prer	medic	ation	IV	2	:0G	220	3	
Did Alleigies																AN	NES	Ť									1	R		L		
																_	JRG	\top			+		\neg				Site			_		
TIME																30	JINO	_			_						SILE	,	_			
Oxygen (LPM)	Т					Т					Т					Т		Т		Т		\top		Т					Т		П	
Nitraus Oxide (LPM)	╈		_	\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+		\dashv	
NIII das Oxide (LPM)	+		_	\vdash		+		_	\vdash		+		\dashv	_		+		+		+		+		\dashv			\vdash		+	Used	-	Waste
Midazolam (mg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+	0360	4	11926
Diazepam (mg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+	—	\dashv	
Diazepam (mg) Fentanyl (mog)	+			\vdash		+		_	\vdash		+		\dashv			+		+		+		+		\dashv	_		\vdash		+	_	\dashv	
Meperidine (mg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+	_	\dashv	
Hydromorphone (mg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+	_	\dashv	
Sufentanil (mcg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		+		\dashv			\vdash		+	—	\dashv	
Ketorolac (mg)	+		_	\vdash		+		_	⊢		+		\dashv	_		+		+		+		+		\dashv			⊢		+	_	\dashv	
Dexamethasone (mg)	+			\vdash		+		_	\vdash		+		\dashv	_		+		+		+		+		\dashv			\vdash		+		\dashv	
NS (mL)	+		_	\vdash		+		_	\vdash		+		\dashv	_		+		+		+		+		\dashv			\vdash		+		\dashv	
0.5% bupivacaine 1:200K epi (mg)	+			\vdash		+			\vdash		+		\dashv			+		+		+		\pm		\dashv			\vdash		+		\dashv	
2% lidocaine 1:100K epi (mg)	+					+		_	\vdash		+		\dashv			+		\pm		+		\pm		_			\vdash		+		\dashv	
4% articaine 1:100K epi (mg)	+			\vdash		†			\vdash		+		\neg			$^{+}$		\top		\top		\top		┪			\vdash		+		┪	
Response to Verbal Stimuli	╈			\vdash		+			\vdash		+		\neg			$^{+}$		+		\top				┪			\vdash		+		┪	
SpO ₂	╅			Г		Ť			Г		\top		\neg			\top		T		1				┪			Т		\top	_	┪	
Respiratory Rate	╅			Г		T			Г		\top		\neg			\top		\top		\top				╛			Г		\top	_	╛	
EtCO₂ Present	十			Т		†			Т		\top		\neg			\top		\top		\top		\top		┪			\vdash		十		┪	
ECG Pattern	╅			Г		\top			Г		\top		\neg			\top		\top		\top		\top		┪			Г		\top	_	\neg	
																						-										
Discharged to:	Τ					Ι					Т					Т	П	Т	П	Т									T	\Box	П	
		\perp	\Box			1	\perp	\perp	Ц	\perp	\perp	\Box			\perp	\perp	Ш	\perp	Ш	\perp	\perp	\perp				\perp	Ц	\perp	\perp	\perp	П	\perp
	L	\perp	Ш			1	\perp	\perp	Ц	_	┸	\perp	Ц	\Box	\perp	┸	Ш	\perp	Ш	4		_	Ш	Ц		\perp	Ц	\perp	\perp	\perp	Ц	\perp
BP	L	4	Ш	Ш		4	4	┺	Ц	4	┸	Ш	Ц	Ц	_	┺	ш	4	\sqcup	4	\perp	_	ш	Ц	_	┺	Ц	4	4	╨	Ц	\perp
15	٦L	_	Ш	Ц	\perp	4	_	┸	Ц	_	┸	Ш	Ц	Ц	_	┸	ш	4	\sqcup	4	ш	_	Ш	Ц	\perp	┸	Ц	_	4	╨	Ц	\bot
HR 15	Ľ	+	Н	Н	-	4	+	╄	Н	4	╄	┺	Ц	4	_	╄	₩	4	+	4	+	4	Н	4	+	╀	Ц	4	4	╨	Н	-
	L	+	Н	Н	-	4	+	╄	Н	4	╀	╄	Н	4	+	╀	₩	4	+	4	\perp	-	Н	4	-	╀	Н	4	+	₩	Н	+
	⊦	+	Н	Н	+	4	+	╀	Н	+	╀	\vdash	Н	\dashv	+	╀	₩	+	₩	4	+	+	Н	-	+	+	Н	+	+	₩	Н	+
SpO ₂	Н	+	Н	Н	-	+	+	╀	Н	+	╀	\vdash	Н	\dashv	+	╀	₩	+	₩	+	+	+	Н	\dashv	+	+	Н	+	+	┯	Н	+
10	0	+	Н	Н	-	+	+	+	Н	+	+	+	Н	+	+	+	₩	+	₩	+	+	+	Н	-	+	+	Н	+	+	₩	Н	+
RR	Н	+	Н	Н	+	+	+	+	Н	+	┿	\vdash	Н	+	+	┿	₩	+	₩	+	+	+	Н	\dashv	+	+	Н	+	+	┯	Н	+
	H	+	\forall	Н	+	+	+	+	H	+	+	\vdash	Н	+	+	+	++	+	+	+	+	+	\forall	\dashv	+	+	Н	+	+	+	$\vdash \vdash$	+
□ Q x 3	H	+	Н	Н	+	+	+	+	H	+	+	\vdash	Н	+	+	+	₩	+	++	+	+	+	Н	\dashv	+	+	Н	+	+	+	\vdash	+
	ŀ	+	Н	Н	+	+	+	+	H	+	+	\vdash	Н	\dashv	+	+	++	+	+	+	+	+	\forall	\dashv	+	+	Н	+	+	+	$\vdash \vdash$	+
☐ Awake 50	ŀ	+	Н	Н	+	+	+	+	H	+	+	\vdash	Н	+	+	+	++	+	++	+	+	+	\forall	┪	+	+	Н	+	+	+	\vdash	+
	H	+	Н	Н	+	+	+	\top	H	+	+	\vdash	Н	\dashv	+	$^{+}$	+	+	+	+	\top	+	Н	\dashv	+	\top	Н	+	+	\top	\vdash	+
■ Alert	h	\top	Н	Н	\top	†	\top	\top	М	\top	\top	т	Н	\dashv	\top	†	+	\top	++	+	\top	\top	Н	\dashv	\top	\top	Н	\top	十	\top	\sqcap	\top
□ Ambulaton.	r	\top	Н			†	+	\top	Н	\top	$^{+}$	П	Н	\dashv	+	$^{+}$	+	\top	+	\top	\top	1	Н	┪	\top	\top	Н	\top	十	\top	\sqcap	
■ Ambulatory	T		П			T	\top	Т	П	\top	Т	П	П	\Box	\top	Τ	\sqcap	\top	\top	1	\top	1	П	┪	\top	Т	П	\top	\top	\top	\sqcap	\top
	٠,	omn	onto			_	_	_	_	_	_		_		_	_		_		_				_		_	_	_	_		_	
	_	Comm	ients	5																												
	-																												_	_		
	-																												_	_		
	-			-		_										-											_		_	_		
Operator Signature														Δ	nesth	Me	nitor	Sion	ature:													



(ofc) 971.801.3394 (cell) 503.339.9568 (fax) extrationpractice@gmail.com www.OregonWisdomTeeth.com

DISCLOSURE AND CONSENT—DENTAL AND ORAL SURGERY

TO THE PATIENT: You have the right, as a patient, to be informed about your condition and about the recommended surgical, medical, or diagnost procedures to be used so that you may make the decision whether or not to undergo the procedure after knowing the risks and hazards involved. The
disclosure is not meant to scare or alarm you; it is simply an effort to make you better informed so you can give or withhold your consent to the procedure.
I voluntarily request Paul A. Brannen, DMD, LLC and such associates, technical assistants, and other healthcare providers as they may deem necessary, to treat my condition which has been explained to me as:
Non-restorable, periodontally-involved, and/or impacted teeth
I(we) understand that the following surgical, medical, and/or diagnostic procedures are planned for me(us), and I(we) voluntarily consent and authorize the procedures under local anesthesia supplemental by: Nitrous Oxide IV Sedation Oral Sedation Local Anesthetic
Surgical extraction of teeth
I(we) understand that my doctor may discover other or different conditions which require additional or different procedures than those planned. I(we) authorize my doctor and such associates, technical assistants, and other health care providers to perform such other procedures which are advisable in their professional judgment.
I(we) understand that no warranty or guarantee has been made to me as to result or cure. I(we) have been given both oral and written post-operative instructions, and I(we) agree to personally contact Dr. Brannen in the event I(we) have a problem. I(we) will follow his instructions until that problem has been satisfactorily resolved. I(we) realize that in the event I(we) develop certain complications, I(we) may miss school or work schedules or I(we) may including, but not limited to, expenses for other dentists, doctors, or medical facilities.
I(we) understand Dr. Brannen is not employed by my dentist but is an independent contractor and will receive a portion of the fee paid to my dentist for these services. I(we) have chosen Dr. Brannen from the alternatives I(we) have been offered to perform my dental surgery. I(we) understand that Dr. Brannen is a General Dentist.
Just as there may be risks and hazards in continuing my present condition without treatment, there are also risks and hazards related to the performance of the surgical, medical, and/or diagnostic procedures planned for me. I(we) realize that common to surgical, medical, and/or diagnostic procedures is the potential for infection, pain, swelling, bleeding, bruising, allergic reactions, and even death. I(we) also realize that the following risks and hazards may occur in connection with this particular procedure:
1. Temporary or permanent nerve injury resulting in altered sensations or numbness of the lips, chin, tongue, teeth, and/or gums.
2. Damage to adjacent teeth and/or dental restorations.
3. Soreness at injection sites and/or along veins, as well as discoloration of the injection sites, face, and/or jaws.
4. Opening of the sinus requiring additional treatment.
5. Jaw fracture, muscle spasms, and/or limited opening of jaws for several days or weeks.
6. Small root fragments remaining in the jaw due to an increased possibility of surgical complications.
7. Jaw joint (TMJ) tenderness, soreness, pain, or locking, which may be temporary or permanent.
8. Other
I(we) understand that IV conscious sedation ("twilight sleep") and other forms of supplemental sedation involve additional risks and hazards, but I(we) request the use of IV conscious sedation and/or other forms of supplemental anesthesia to assist in the relief and protection from pain during the planned an additional procedures. I(we) realize the IV conscious sedation and/or other forms of supplemental anesthesia may have to be changed possibly without explanation to me(us). I(we) understand this is not general anesthesia (being completely asleep), and that it is unlikely, but I may have unpleasant memorie of the procedure.
I(we) understand that certain complications may result from the use of any IV sedative or other form of anesthesia, including respiratory problems, drug reactions, paralysis, brain damage, or even death. Other risks and hazards which may result from the use of IV sedation or other sedatives or anesthetics range from minor discomfort to injury of the vocal chords, teeth, and/or eyes.
I(we) have been given an opportunity to ask questions about my(our) condition, alternative forms of anesthesia and treatment, risks of non-treatment, the procedures to be used, and the risks and hazards involved, and I(we) believe that I(we) have sufficient information to give this consent.
I(we) certify this form has been fully explained to me(us), that I(we) have read it or have had it read to me(us), that the blank spaces have been filled in, and that I(we) understand its contents
DATE: TIME:
Signature of Patient or Other Legally-responsible Person / Patient's Name (Please Print)
WITNESS: DATE:



(ofc)	971.801.3394 (cell)	503.339.9568 (fax)	extrationpractice@gmail.com
	www.Orego	nWisdomTeeth.com	

POST-OPERATIVE INSTRUCTIONS FOR THE SEDATION PATIENT

- Patients may sleep but must be watched for at least six (6) hours after treatment.
- Please escort the patient from the vehicle to the house by supporting them under the arm. Do not let the patient go up or down stairs unescorted.
- Patient may not drive for 24 hours after being sedated and must limit physical activity.
- Give the patient medications only as directed by Dr. Brannen.
- Leave the "Post-operative Instructions" form and any medication easily available for the patient.

PAIN MANAGEMENT

Ibuprofen = Advil, Motrin
—600mg (3 tablets of 200mg) Ibuprofen taken every 6 hours—
NEVER take more than 3200 mg (16 x 200mg tablets) Ibuprofen in 24 hours.

AND

Acetaminophen = Tylenol

—500-1000 mg Acetaminophen (1-2 tablets of 500mg Tylenol) every 6 hours— NEVER take more than 4000 mg (8 x 500mg tablets) Acetaminophen in 24 hours.

If you have a prescription for Vicodin or Percocet, it will be substituted in place of Acetaminophen

If you have any questions about this regimen or about the post-operative instructions, please call:

Paul Brannen, DMD, LLC 971.801.3394

RELEASE

I am escorting (patient's name)	home. I understand that the patient
has been sedated. I have received a copy of the posagree to follow these instructions. I have had an oppor	st-operative instructions for the sedated patient, and I
Escort's Name (please print)	
Escort's Signature	Date
Witness's Name (Doctor/Assistant—please print)	
Witness' Signature	Date



(ofc) 971.801.3394 (cell) 503.339.9568 (fax) extrationpractice@gmail.com www.OregonWisdomTeeth.com

POST-OPERATIVE INSTRUCTIONS FOLLOWING DENTAL SURGERY

THINGS TO EXPECT:

Bleeding: Bleeding or "oozing" for the first 12 to 24 hours.

<u>Swelling</u>: This is normal following a surgical procedure in the mouth. It should reach its maximum in two-to-three days

and should begin to diminish by the fifth post-operative day.

Discomfort: The most discomfort that you may experience may occur for a few hours after the sensation returns to your

mouth. It may gradually increase again for 2-3 days, then begin to diminish over the next few days.

THINGS TO DO IMMEDIATELY FOLLOWING SURGERY:

<u>Bleeding:</u> Place gauze over extraction sites and maintain pressure by biting for at least one hour. Repeat as needed. Keep

head elevated, and rest. Do not suck or spit excessively. (Also, please refrain from blowing into musical

instruments.)

NOTE: Some "oozing" and discoloration of saliva is normal. If bleeding persists, replace the gauze with a clean

folded gauze placed over the extraction site, and maintain the pressure until the bleeding stops.

Swelling: Place ice or cold compresses on the region of surgery for ten minutes every half-hour for the first eight to 12

hours.

NOTE: Ice bags or cold compresses should be used only on the day of surgery.

Smoking: Avoid smoking during the healing period.

Discomfort: Take medications as directed for <u>PAIN</u>. Mild-to-moderate pain can be relieved by non-prescription Advil,

Aleve, or Orudis. For more severe pain, take the prescription pain medication as directed. Remember that these medications can take up to 30 minutes to one hour to take effect. If you are using any of these medications for

the first time, exercise caution with the initial doses (start with ½ a pill).

<u>Diet</u>: A nutritious liquid or soft diet will be necessary for the first weeks after surgery. Healing will occur in weekly

increments; therefore, it is best to gradually (in weekly increments) return the diet and/or other mouth/oral

activities back to normal.

Physical For the first 24 to 48 hours, one should REST. Patients who have sedation should refrain

<u>Activity</u>: from driving an automobile or from engaging in any task that requires alertness for the next 24 hours.

THE DAYS AFTER SURGERY:

- 1. Brush teeth carefully.
- 2. Beginning 24 hours after the surgery, rinse mouth with <u>WARM SALT WATER</u> (or prescription mouth rinse). Continue rinsing three-to-five times per day for seven days, then begin irrigating per dentist's instructions (see #7 below).
- 3. If <u>ANTIBIOTICS</u> are prescribed, be <u>SURE</u> to take <u>ALL</u> that have been prescribed, <u>AS DIRECTED</u>.
- 4. Use <u>WARM, MOIST HEAT</u> on face for swelling, if any. Continue until the swelling subsides. A warm, wet washcloth or heating pad will suffice.
- 5. If SUTURES were used, they will dissolve on their own.
- 6. <u>DRY SOCKET</u> is a delayed healing response, which may occur during the second to fourth post-operative day. It is associated with a throbbing pain on the side of the face, which may seem to be directed up toward the ear. In mild cases, simply increasing the pain medication can control the symptoms. If this is unsuccessful, please call Dr. Brannen.
- 7. RETURN TO YOUR DENTIST'S OFFICE five-to-seven days after the surgery for irrigation instructions.

CONTACT THE DOCTOR IF:

- 1. Bleeding is excessive and cannot be controlled.
- 2. Discomfort is poorly controlled.
- 3. Swelling is excessive, spreading, or continuing to enlarge after 60 hours.
- 4. Allergic reactions to medications occur, which are causing a generalized rash or excessive itching.

CONTACT EMERGENCY MEDICAL SERVICES ("EMS") OR CALL "911" IF:

Patient loses or has lost consciousness.



(ofc)	971.801.3394 (cell)	503.339.9568 (fax)	extrationpractice@gmail.com
		www.OregonWisdo	mTeeth.com

ACKNOWLEDGMENT: RECEIPT OF NOTICE OF PRIVACY PRACTICES

I have received a copy of Paul A. Brannen, D	OMD, LLC's Notice of Privacy Practices effective 3/1/17.
Patient's Name (please print)	
Signature of Patient	Date Signed
*******	*********
I am a parent or legal guardian of received a copy of Paul A. Brannen, DMD, L	(patient's name). I have LC's Notice of Privacy Practices effective 3/1/17.
Parent or Legal Guardian's Name (please prin	nt)
Relationship to Patient: Parent	Legal Guardian
Signature of Parent or Legal Guardian	Date Signed
I authorize the doctor and his staff to contact	me byphoneemailmail (check all that apply)
*******	*******
	rdian did not sign above, staff member must document when al, why the acknowledgment could not be obtained, and what
Notice of Privacy Practices effective 3/1/17 g	given to individual on (date)
☐ In Person ☐ Email ☐ Mail ☐ Othe	er
Reason patient or patient's parent/legal guard	lian did not sign this form:
Did not want to sign Did not respond after more than one atten Other	npt
Staff Member's Name (please print)	Title
Signature of Staff Member	Date Signed